

Trinity College Interlibrary Loan

ILLiad TN: 71653



Borrower: COF

Lending String: *TYC,TXA,QPE,GZM,EYM

Patron: Nemeth, Nicole/MICRO/g/DELIVER

Journal Title: Cotinga.

Volume: 21 **Issue:** .
Month/Year: 2004 **Pages:** .

Article Author:

Article Title: 'Nemeth, N. and C. Vaughan "Daily observations of Scarlet Macaws (Ara macao) in Costa Rica"'

Imprint: Sandy, Bedfordshire, U.K. ; Neotropical

ILL Number: 6405640



Call #: current QL671 .C68

Location: Watkinson

ARIEL
Charge
Maxcost: 50ifm

Shipping Address:
Colorado State University
Libraries/ILL
200 W Lake St.
Fort Collins, CO 80523-0002

Fax: 970-491-2252
Ariel: 129.82.31.241

F. W. 13

Charge

lighting. The bird's size excludes all New World swifts^{5,7}, except the largest, White-collared Swift *Streptoprocne zonaris* which, however, possesses quite different, largely deep black plumage⁷. The only species to obviously accord with my description is Alpine Swift⁵, with which I have experience from one observation in India. There have been four records of this species in the Caribbean³: one on 20 September 1955 in Barbados¹, which was collected; one on 14 April 1987 on Guadeloupe⁴; one on 20 July 1987 on Desecheo Island, Puerto Rico⁵ and one on 19 August 1992 on St Lucia², which was photographed. Mottled Swift *Tachymarptis aequatorialis*, the only congeneric, has a longer, more deeply forked tail and lacks the large white oval belly patch. Furthermore, the latter has not been reported from the Americas⁵.

References

1. Bond, J. (1985) *The birds of the West Indies*. Fifth edition. London, UK: Collins.
2. Burke, W. (1994) Alpine Swift (*Tachymarptis melba*) photographed on St. Lucia, Lesser Antilles—third record for the Western Hemisphere. *El Pitirre* 7 (3): 3.

3. Ebels, E. B. (2002) Transatlantic vagrancy of Palearctic species to the Caribbean region. *Dutch Birding* 24: 202-209.
4. Feldman, P. & Pavis, C. (1995) Alpine Swift (*Tachymarptis melba*) observed in Guadeloupe, Lesser Antilles: a fourth record for the Western Hemisphere. *El Pitirre* 8 (2): 2.
5. del Hoyo, J., Elliott, A. & Sargatal, J. (1999) *Handbook of the birds of the world*, 5. Barcelona: Lynx Edicions.
6. Meier, A. J., Noble, R. E. & Raffaele, H. A. (1989) The birds of Desecheo Island, Puerto Rico, including a new record for Puerto Rican territory. *Carib. J. Sci.* 25: 24-29.
7. Meyer de Schauensee, R. & Phelps, W. H. (1978) *A guide to the birds of Venezuela*. Princeton, NJ: Princeton University Press.

Otte Ottema

Bird Department, Stinasu Foundation for Nature Conservation in Suriname, Cornelis Jongbawstraat 14, P.O. Box 12252, Paramaribo, Surinam. E-mail: research@stinastu.sr.

Feeding observations on Scarlet Macaw *Ara macao* in Costa Rica

Nicole M. Nemeth and Christopher Vaughan

Cotinga 21 (2004): 71-72

Observamos un grupo familiar de cuatro individuos, dos adultos y dos jóvenes, de Lapa Roja *Ara macao* alimentándose de los frutos de dos especies de árboles. Los adultos consumían los frutos de la Palma Royal *Scheelea rostrata*. Los dos juveniles, aparentemente carentes de habilidad para alimentarse de éstos, consumían al mismo tiempo los frutos del Jobo *Spondias mombin*. La disponibilidad de diferentes recursos alimentarios puede incrementar la supervivencia de juveniles, los cuales aún no han perfeccionado sus diferentes técnicas de alimentación.

On 7 September 1995, from approximately 14h00-14h15, four Scarlet Macaws *Ara macao* were observed feeding near the Carara National Park guard station, Costa Rica (09°47'N 84°35'W). They were feeding on the edge of partially deforested pasture, along the Costañera highway. Two adults and their two juveniles were concurrently feeding on two different tree species, both common in the region. The two adults were eating from a Royal Palm *Scheelea rostrata*, of approximately 30-35 m height, which contained large bunches of mature palm nuts. *S. rostrata* has been observed with abundant nut clusters at various times throughout the year, and is also common in seasonally inundated areas in central Pacific Costa Rica (pers. obs.)². *Scheelea rostrata* is a well-known, abundant food source for *A. macao* in the region, and is high in nutritional value^{2,3}. The nuts hang inverted from the

top of the tree, and must be removed by *A. macao* in mid-air. Within the same flight, the nuts are carried to a nearby tree for consumption. Therefore, the removal of palm nuts requires advanced skills, probably still under-developed in the two juveniles (estimated at 3-4 months old).

The juveniles were feeding c.20 m off the ground in a nearby Jobo *Spondias mombin*, which provided many perches as well as nuts surrounded by visible, yellow fruits³. The base of the *Spondias mombin* tree was within 10-20 m of the *Scheelea rostrata*, and all four birds vocalised throughout the observation; the juveniles emitted soft, single-syllable calls while the adults shrieked loudly. We have observed both juvenile and adult *A. macao* consuming *Spondias mombin* at various times throughout the year. The adults flew repeatedly to the *Scheelea rostrata* to remove fruits swiftly from the tree,

immediately landing in nearby trees to consume portions of the nuts before intentionally dropping them. Simultaneously, the juveniles remained in the same *Spondias mombin*, walking among its branches in an uncoordinated fashion, often appearing to lose their balance, resulting in dropped fruits.

Although it has been suggested that skill variation among conspecifics, including juveniles versus adults, may affect survival¹, there appears to be a variety of food sources available to *A. macao* throughout the year (CV unpubl. data). These available food sources would enhance the survival rate of juveniles, which have not yet perfected various feeding techniques. Further, it has been suggested that adult *A. macao* lack the strength to access all parts of the palm fruit, and *Scheelea rostrata* may not be an essential part of their diet (S. Matola pers. comm.). However, in our study, *Scheelea rostrata* comprised a significant portion of feeding observations (38 of 163, or 23.3%), which included 43 different plant species (CV unpubl. data). Little has been published about the Central Pacific population of *A. macao*, which is one of two significant populations remaining in Costa Rica^{4,5}.

Acknowledgements

We would like to thank The Organization of American States, the Universidad Nacional de Costa Rica, Idea Wild and Wildlife Trust for financial support, and the guards at Carara National Park for housing during the study. Fernando González García helped prepare the manuscript.

References

1. Garnett, S. & Crowley, G. (1995) Feeding ecology of hooded parrots *Psephotus dissimilis* during the early wet season. *Emu* 95: 54–61.
2. Janzen, D. H. (1975) *Ecology of plants in the tropics*. London, UK: Edward Arnold.
3. Janzen, D. H. (1983) *Costa Rican natural history*. Chicago: University of Chicago Press.
4. Marineros, L. & Vaughan, C. (1995) Scarlet Macaws of Carara. In: Abramson, J., Speer, B. & Thomsen, J. (eds.) *The large macaws: their care, breeding and conservation*. California: Raintree Publications.
5. Vaughan, C., McCoy, M. & Liske, J. (1991) Ecology and management of the Scarlet Macaw in Carara Biological Reserve. In: Clinton-Eitnicar, J. (ed.) *Proc. First Mesoamerican workshop on the conservation and management of macaws*. San Antonio, TX: Center for the Study of Tropical Birds.

Nicole M. Nemeth and Christopher Vaughan

Regional Wildlife Management Program for Mesoamerica and the Caribbean, Universidad Nacional de Costa Rica, Apartado 1350-3000, Heredia, Costa Rica. NMN currently at: College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Ft. Collins, CO 80523, USA. E-mail: nnemeth@colostate.edu and CV currently at: Institute for Environmental Studies, University of Wisconsin, Madison, WI 53706, USA. E-mail: cvaughan@facstaff.wisc.edu.

Primer registro de *Arenaria interpres* para Bolivia

Mauricio Herrera

Cotinga 21 (2004): 72–73

Ruddy Turnstone *Arenaria interpres*, a boreal migrant in South America, is abundant along the coasts of Peru to Chile. During research with WWF-Bolivia, I observed two individuals in a mixed flock of shorebirds at Lake Cáceres, prov. Germán Bush, dpto. Santa Cruz, on 26–28 October 1998, accompanied by Spotted Sandpiper *Actitis macularia* and American Golden Plover *Pluvialis dominica*. This is the first record of *A. interpres* for Bolivia. I collected a juvenile male, which has been deposited in the Noel Kempff Mercado Natural History Museum (MNKM 1692). Lake Cáceres is near Puerto Suárez, 645 km south-east of the city of Santa Cruz. There are few records of Ruddy Turnstone from the interior of South America.

Arenaria interpres es un playero de distribución cosmopolita. Es un migrante boreal en Sudamérica, abundante sólo por las costas marinas en el norte del continente, principalmente Perú y el norte y centro de Chile, siendo escaso pero predecible en Argentina¹.

Durante el trabajo de campo del proyecto Evaluación y Identificación de Ecosistemas y

Especies Importantes para la Conservación en las Áreas Protegidas del Pantanal, financiado por la WWF-Bolivia, se observaron dos individuos de *Arenaria interpres* en un grupo mixto de aves playeras, alimentándose en las orillas de la laguna Cáceres, Estancia Arco Iris, prov. Germán Bush, dpto. Santa Cruz (18°56'S 57°48'W). Fueron observados sólo por las mañanas de los días 26–28